

The black hole fundamental plane: revisit with a larger sample of radio and X-ray emitting broad-line AGNs

Zhao-Yu Li, Xue-Bing Wu and Ran Wang

Department of Astronomy, Peking University, Beijing 100871, China

zhaoyuli@pku.edu.cn; wuxb@bac.pku.edu.cn; littlestar@pku.edu.cn

ABSTRACT

We use a recently released SDSS catalog of X-ray emitting AGNs in conjunction with the FIRST 20cm radio survey to investigate the black hole fundamental plane relationship between the 1.4GHz radio luminosity (L_r), 0.1-2.4 keV X-ray luminosity (L_X), and the black hole mass (M), namely, $\log L_r = \xi_{RX} \log L_X + \xi_{RM} \log M + \text{const.}$ For this purpose, we have compiled a large sample of 725 broad-line AGNs, which consists of 498 radio-loud sources and 227 radio-quiet sources. Our results are generally consistent with those in our previous work based on a smaller sample of 115 SDSS AGNs. We confirm that radio-loud objects have a steeper slope (ξ_{RX}) in the radio-X-ray relationship with respect to radio-quiet objects, and the dependence of the black hole fundamental plane on the black hole mass (ξ_{RM}) is weak. We also find a tight correlation with a similar slope between the soft X-ray luminosity and broad emission line luminosity for both radio-loud and radio-quiet AGNs, which implies that their soft X-ray emission is unbeamed and probably related to the accretion process. With the current larger sample of AGNs, we are able to study the redshift evolution of the black hole fundamental plane relation for both radio-loud and radio-quiet subsamples. We find that there is no clear evidence of evolution for radio-quiet AGNs, while for radio-loud ones there is a weak trend where ξ_{RM} decreases as the redshift increases. This may be understood in part as due to the observed evolution of the radio spectral index as a function of redshift. Finally, we discuss the relativistic beaming effect and some other uncertainties related to the black hole fundamental plane. We conclude that, although introducing scatters to the fundamental plane relation, Doppler boosting alone is not enough to explain the observed steeper value of ξ_{RX} in the radio-loud subsample with respect to the radio-quiet ones. Therefore, the significant difference of ξ_{RX} between radio-loud and radio-quiet sources is probably also due to the different physical properties of the jets.

Subject headings: accretion, accretion disks – black hole physics – galaxies: active
 – galaxies: nuclei – radio continuum: galaxies – X-ray: galaxies

1. INTRODUCTION

Astrophysical black holes do not emit light directly, but they can be probed by their gravitational influence on neighboring matter, which produces observable signatures of black hole activity. The key mechanism for a black hole to become active is the accretion process (Frank, King & Raine 2002), which is usually accompanied by a relativistic jet. Accretion disks and jets can produce photons from radio to X-ray band. The radio emission is usually believed to originate from the synchrotron radiation of the jet (Begelman, Blandford & Rees 1984), while the optical/UV emission mostly comes from the multicolor black body radiation emitted from the accretion disk (Shakura & Sunyaev 1973), and the X-ray radiation is usually associated with the inner-most region of the accretion disk, where the temperature is the highest. In some cases, the contribution of inverse Compton scattering from high energy electrons in a disk corona is also needed to account for the observed power-law spectrum in the X-ray band (Haardt & Maraschi 1993). If jet production is directly related to the accretion process, we would expect a natural correlation between radio and X-ray luminosities (Merloni, Heinz & Di Matteo 2003; Heinz & Sunyaev 2003; Falcke et al. 2004).

The radio-to-X-ray correlation has long been studied in both Galactic black hole (GBH) candidates and active galactic nuclei (AGNs). Gallo, Fender & Pooley (2003) found a strong correlation between the radio and X-ray emission ($L_r \propto L_X^{0.7}$) using the simultaneous X-ray and radio observational data of stellar-mass black hole X-ray binaries (XRBs) during the low/hard state. In addition, they suggested that when XRBs enter the hard to soft transition state, the jet is suppressed and the radio emission decreases. Recently, some studies have shown that the substantial scatter exists in such a relationship of GBHs (Gallo 2006; Xue & Cui 2007; Xue, Wu & Cui 2008). In the case of AGNs, Brinkmann, Yuan & Siebert (1997) obtained a remarkable correlation between 2keV X-ray luminosity and 5GHz radio luminosity for 324 radio-loud AGNs. Canosa et al. (1999) found a strong correlation between soft X-ray and 5GHz radio luminosities for 40 low-power radio galaxies. Brinkmann et al. (2000) studied a sample derived from the cross-correlation of the *ROSAT* All-Sky Survey (RASS) catalog and the Very Large Array (VLA) Faint Images of the Radio Sky at Twenty-Centimeters (FIRST 20cm) catalog. They found that for 843 AGNs, the X-ray and radio luminosities are correlated over two decades in radio luminosity, spanning radio-loud and radio-quiet regimes, but radio-quiet quasars seem to follow a different correlation from

radio-loud ones. Recently, Panessa et al. (2007) investigated the radio/X-ray luminosity correlation for low-luminosity AGNs, including local Seyfert galaxies and low luminosity radio galaxies (LLRGs). They found that X-ray and radio luminosities are significantly correlated over 8 orders of magnitude, with Seyfert galaxies and LLRGs showing the similar slope, which seems different from the previous results.

Theoretical explanations for the observed radio-to-X-ray relation have been discussed in a number of previous works. Fender et al. (2003) found that, at relatively lower accretion rate ($\dot{M} < 7 \times 10^{-5} \dot{M}_{Edd}$), black hole X-ray binaries would enter a “jet-dominated” state. At this stage, the majority of the liberated accretion power is transferred into the jet and does not dissipate as X-ray emission in the accretion flow. This was also suggested by Falcke, K rding & Markoff (2004), who demonstrated that, below a critical value of the accretion rate ($< 1 \sim 10\% \dot{M}_{Edd}$), the Spectral Energy Distribution (SED) of a black hole accreting source is dominated by the non-thermal radiation from the jet, while for sources with higher accretion rates ($\dot{M} \leq \dot{M}_{Edd}$), the radiation is dominated by the accretion flow. Using a coupled disk-jet model, Yuan & Cui (2005) showed that, in GBHs when the X-ray luminosity is smaller than a certain value ($\sim < 10^{-5} - 10^{-6} L_{Edd}$), the jet will dominate the radiation and the radio-to-X-ray correlation becomes to $L_r \propto L_X^{1.23}$. When the X-ray luminosity exceeds that critical value, the accretion flow would produce most of the X-ray emission and the radio-to-X-ray correlation then becomes to $L_r \propto L_X^{0.7}$. Therefore, the accretion rate decides the detailed physical model of the accretion disk, which then leads to different observed radio-to-X-ray correlation slopes (K rding, Falcke & Corbel 2006). Although it is still not clear whether such a critical accretion rate is of the same order of magnitude for both GBHs and AGNs, the similarity of accretion-jet physics in these two systems seems to imply that similar results may also be found in AGNs.

In a comprehensive study, Merloni et al. (2003) examined a sample combining Galactic black hole systems and supermassive black hole systems by investigating their compact emission in X-ray (2-10 keV) and radio (5 GHz) bands. They found that the radio luminosity (L_r) is strongly correlated with both the black hole mass (M) and the X-ray luminosity (L_X) (so called the fundamental plane of black hole activity). The relation is: $\log L_r = (0.60^{+0.11}_{-0.11}) \log L_X + (0.78^{+0.01}_{-0.09}) \log M + (7.33^{+4.05}_{-4.07})$. Subsequently, Wang et al. (2006) selected a uniform sample of broad-line AGNs which was cross-identified from the RASS, Sloan Digital Sky Survey (SDSS) and FIRST 20cm radio survey to test the black hole fundamental plane relation. Their final sample consisted of 115 broad emission line AGNs including 39 radio-quiet AGNs and 76 radio-loud ones. They found that the relationship has a very weak dependence on the black hole mass. Moreover, radio-quiet and radio-loud objects have different radio-to-X-ray slopes, which is 0.85 for radio-quiet objects and 1.39 for radio-loud sources. This differs from the result of Merloni et al. (2003) where the relationship seems

to be universal for different types of black hole sources. However, the limited statistics of the sample in Wang et al. (2006) motivated us to increase the number of sources in order to confirm the results on stronger statistical bases. With a larger sample, we are also able to study the possible evolution of the black hole fundamental plane relation as we have enough sources in each redshift bins.

We organize the paper as the following. In §2 we present our sample selection criteria and the main properties of our sample. In §3 we show the derived fundamental plane relation and investigate the possible evolution of such a relation. In §4 we briefly discuss and summarize our results.

2. THE RADIO AND X-RAY EMITTING BROAD-LINE AGN SAMPLE

2.1. Sample selection

Our sample is selected based on the cross-identification of the newly published X-ray emitting SDSS AGN catalog (Anderson et al. 2007) and the catalog of the FIRST 20cm radio survey¹ (White et al. 1997). Anderson et al. (2007) employed X-ray data from the *ROSAT* All-Sky Survey (RASS) and both optical imaging and spectroscopic data from the Sloan Digital Sky Survey (SDSS). It is worth emphasizing that the RASS and SDSS are extremely well matched to each other via a variety of coincidences (e.g., similar survey depth, sensitivity and sky coverage, etc.)(Anderson et al. 2003). The RASS/SDSS data from 5740 deg² of sky spectroscopically covered in SDSS Data Release 5 provide an expanded catalog of 7000 confirmed quasars and other AGNs that are probable RASS identifications (Anderson et al. 2007).

One of the main benefits of this sample is that the SDSS surveyed area is also covered by the FIRST 20cm radio survey. We cross-correlate the broad-line AGN catalog of Anderson et al. (2007) with the FIRST radio-detected sources to build a RASS-SDSS-FIRST cross-identified sample of 868 broad-line AGNs. All of these 868 sources have been observed and detected at 1.4GHz (FIRST 20cm survey) and in 0.1-2.4 keV energy band (RASS). The optical spectra from the SDSS data archive can be used to estimate the central black hole mass. Here, we exclude the high z sources ($z > 2.171$) in our sample for the lack of the $H\beta$ and $Mg\ II\ \lambda 2798$ emission lines in their SDSS optical spectra. Black hole masses of the remaining 725 sources are estimated. The radio-loudness (R) is calculated with the rest-frame B-band (4400Å) and 5GHz flux density according to the definition $R = f_{5GHz}/f_B$

¹See VizieR Online Data Catalog, 8071 (Becker et al. 2003)

(Kellermann et al. 1989), and the 5GHz flux density is derived from the 1.4GHz flux density assuming a spectral index of 0.5 (Kellermann et al. 1989). Radio-loud and radio-quiet sources are separated by $R = 10$. Please see subsection 2.2 of Wang et al. (2006) for the details of data reduction and black hole mass estimation (Kaspi et al. 2000, 2005; McLure & Jarvis 2002; Wu et al. 2004).

Our final sample comprises 725 entries, of which 498 are radio-loud, 227 are radio-quiet. In addition, we discover several radio-loud narrow-line Seyfert 1 galaxies (NLS1) in our study. A detailed discussion of these radio-loud NLS1s is presented in Appendix A. Throughout this paper, we use the cosmology² with $H_0 = 70 \text{ kms}^{-1}\text{Mpc}^{-1}$, $\Omega_\Lambda = 0.7$ and $\Omega_M = 0.3$.

2.2. Sample properties

We describe here the main properties of our selected AGN sample. Table 1 gives the total 725 sources with the SDSS optical name, redshift, broad-band (0.1-2.4 keV) soft X-ray luminosity, rest-frame 1.4GHz radio luminosity, radio-loudness, black hole mass, broad emission line luminosity and corresponding flag. In Figure 1, histograms of the redshift, logarithm of radio-loudness, 0.1-2.4 keV X-ray luminosity, black hole mass (in unit of $M_\odot$), 1.4GHz radio luminosity and ratio of X-ray to Eddington luminosity (L_{edd})³ of our sample are plotted. The radio-loudness distribution does not show a clear dichotomy between radio-loud and radio-quiet AGNs. This is not surprised for samples of FIRST detected quasars. The radio-loudness dichotomy was found mostly for optically selected samples, and is usually absent for samples of deep radio-detected sources (Brinkmann et al. 2000; White et al. 2000; Lacy et al. 2001).

Our sample is a part of the broad-line AGN catalog provided by Anderson et al. (2007), which includes typical broad-line quasars, Seyfert 1 galaxies, low redshift Seyfert 1.5-1.9 galaxies and some rare galaxies like NLS1s. Compared with Wang et al.(2006), the distribution ranges of different physical parameters in our sample are relatively larger. The significant range in the luminosity at each redshift bin can avoid the strong dependence of luminosity on redshift in a flux-limited sample (Avni & Tananbaum 1982). In particular,

²We now use this new cosmology to calculate the black hole mass from the $H\beta$ broad component's luminosity and FWHM. That is, $M_{bh} = 1.464 \times 10^5 \left(\frac{R_{BLR}}{1 \text{ lt days}} \right) \left(\frac{V_{FWHM}}{10^3 \text{ km/s}} \right)^2$ and $\text{Log} R_{BLR} = (1.324 \pm 0.086) + (0.667 \pm 0.101) \text{Log} \left(\frac{L_{H\beta}}{10^{42} \text{ ergs s}^{-1}} \right)$. The latter relation is an updated version of the $R_{BLR} - L_{H\beta}$ relation proposed by Wu et al. (2004) using the new cosmology.

³Defined as $L_{Edd} = 1.26 \times 10^{38} (M/M_\odot) \text{ ergs s}^{-1}$

more convincing results can be obtained with the larger sample.

However, we must notice the selection biases in our sample. First of all, AGNs with radio flux fainter than the FIRST detection limit can not be detected by FIRST. Therefore, our sample does not include radio-quiet AGNs with $\log_{10}(\text{Radio} - \text{Loudness}) < -0.23$.⁴ In addition, based on the redshift distribution shown in Figure 1, it is clear that most AGNs at high redshifts in our sample are radio-loud. Therefore, any conclusion based on the total sample is biased toward luminous radio-loud AGNs. We also perform the Kolmogorov-Smirnov (K-S) test to evaluate the distribution similarity of physical parameters shown in Figure 1 between radio-quiet and radio-loud subsamples (such as L_r , L_X , M_{BH} , and $\frac{L_X}{L_{\text{Edd}}}$). We found that the probabilities of similarity are all less than 0.05, meaning that the distribution of these physical quantities in the two subsamples are essentially different, with relatively larger mean value of each parameter for radio-loud AGNs.

We plot the radio vs. X-ray luminosity (left panel) and the Eddington-luminosity-scaled radio vs. scaled X-ray luminosity (right panel) in Figure 2. Objects in different black hole mass bins are presented with different symbols and colors to highlight a possible segregation in the plot. Clearly, there is no trend that sources in different black hole mass bins are parallel to each other, which is consistent with the result in Wang et al. (2006). Again, tight correlation between the radio and X-ray luminosities is clear, with radio luminosity spanning more than 6 orders of magnitude.

Figure 3 and Figure 2 are identical except that in Figure 3, different colors and symbols are used to denote different radio-loudness bins instead of black hole mass bins. AGNs in different radio-loudness bins seem to distribute in parallel sequences. The result also confirms the conclusion of Wang et al. (2006).

3. CORRELATION ANALYSIS

In this section, we derive the black hole fundamental plane relation based on the current large sample. We also adopt several statistical methods to test its significance. Finally, we examine the possible evolution of this relation at different redshifts.

⁴ $10^{-0.23}$ is the minimum radio-loudness value of radio-quiet AGNs in our sample

3.1. The fundamental plane relation

In the three-dimensional space ($\log L_r$, $\log L_X$, $\log M$), black hole systems are preferentially distributed on a plane, called the “fundamental plane of black hole activity” (Merloni et al. 2003). Based on our sample of AGNs, we fit the data in the form:

$$\log \left(\frac{L_r}{10^{40} \text{ergs s}^{-1}} \right) = \xi_{RX} \log \left(\frac{L_X}{10^{44} \text{ergs s}^{-1}} \right) + \xi_{RM} \log \left(\frac{M}{10^8 M_\odot} \right) + \text{const.} \quad (1)$$

We also fit the relation between Eddington-luminosity-scaled radio and X-ray luminosities for the radio-quiet subsample only. The fitting formula is

$$\log \left(\frac{L_r}{L_{\text{Edd}}} \right) = \xi_{\text{ERX}} \log \left(\frac{L_X}{L_{\text{Edd}}} \right) + \text{constant.} \quad (2)$$

We apply the Ordinary Least Squares (OLS) multivariate regression method (Isobe et al. 1990) to the total, radio-loud and radio-quiet subsamples, respectively. The OLS bisector fitting result for equation (1) with errors at the one-sigma confidence level and the dispersion (σ_r)⁵ are given in Table 2. We also list results of previous works for comparison. Our result is generally consistent with Wang et al. (2006), but now with smaller uncertainty for each coefficient due to the larger sample. The conclusions are similar: first, the coefficient ξ_{RX} tends to be larger as the radio-loudness increases; second, the black hole mass seems to be unimportant in the correlation between radio and soft X-ray luminosities. We should note that the black hole fundamental plane coefficients calculated by Merloni et al. (2003) are based on a radio-loud and radio-quiet combined sample, with a predominance of radio-quiet sources. Our result for the radio-quiet subsample is similar to that of Merloni et al. (2003) but the dependence on the black hole mass is weaker in our case.

The fitting result for equation (2) with errors at the one-sigma confidence level is

$$\log \left(\frac{L_r}{L_{\text{Edd}}} \right) = (0.96 \pm 0.04) \log \left(\frac{L_X}{L_{\text{Edd}}} \right) + (-4.82 \pm 0.08). \quad (3)$$

The dependence of L_r/L_X on the black hole mass is shown in Figure 4. The overall correlation is weak, except that a positive correlation is observed for sources (mostly radio-quiet ones) with black hole masses smaller than $10^{7.5} M_\odot$. Figure 5 shows the edge-on black hole fundamental plane relation for radio-quiet sources, with radio-loud AGNs overplotted for comparison. We also plot L_r/L_{Edd} vs. L_X/L_{Edd} in Figure 6, with different symbols representing different radio-loudness bins. Parallel sequences can be seen clearly from these figures.

⁵We define the dispersion as the square root of the variance of the differences between the observed radio luminosity and that calculated from the fitting relation

3.2. Statistical tests for the fundamental plane relation

3.2.1. Partial correlation tests

As Bregman et al. (2005) pointed out, the correlation between X-ray and radio luminosities may be dominated by the distance effect. Following Wang et al. (2006), we performed the partial Kendall τ correlation test to examine this effect (Akritas & Siebert 1996). In Table 3, Column (1)-(3) list the variable names of X, Y and Z respectively, where the partial correlation of X and Y is calculated with the influence of Z variable excluded. Column (4) gives the subsample type. Column (5) lists the number of sources in each subsample. Column (6)-(8) show results of the partial correlation test, the square root of the calculation variance and the probability of the null hypothesis. The null hypothesis is rejected with a probability less than the significance level (i.e., ~ 0.05). From Table 3 we can see that the partial correlation between L_r and L_X is strong even after excluding the distance effect, because the P_{null} value is less than 10^{-10} .

3.2.2. The scrambling test

Besides the partial correlation test performed above, we adopt another method introduced by Bregman (2005) to evaluate the degree of influence that any distance effect has on our sample (i.e., Merloni et al. 2006). We calculate the Pearson correlation coefficient (ρ) between $\text{Log} \left(\frac{L_r}{10^{40} \text{ ergs s}^{-1}} \right)$ and $\xi_{RX} \text{Log} \left(\frac{L_X}{10^{44} \text{ ergs s}^{-1}} \right) + \xi_{RM} \text{Log} \left(\frac{M_{bh}}{10^8 M_\odot} \right)$ by randomly assigning radio fluxes to objects in our sample. This procedure is performed 10^6 times in order to construct the Monte Carlo test. The result is shown in Figure 7. The correlation coefficients are adopted from Table 2. For comparison, we overplot the Pearson correlation coefficient of the original sample with a vertical line in each panel of Figure 7.

For radio-quiet objects, our 10^6 realizations of randomized datasets produce only one case, where the Pearson correlation coefficient exceeds the value of the real datasets. For radio-loud sources, *not even one* shows a stronger correlation than the real value. This means that, for the radio-quiet subsample, the probability that the correlation of the fundamental plane is entirely due to the distance effect is about 1×10^{-6} , and it is even less than 10^{-6} for the radio-loud subsample and the entire sample. Therefore, we are confident to say that the existence of a correlation between the radio luminosity, X-ray luminosity and black hole mass cannot be purely the result of distance effects. This result is consistent with the partial correlation test performed above.

3.3. The correlation between soft X-ray and broad emission line luminosities

In Figure 8, we plot the soft X-ray luminosity versus the $H\beta$ broad emission line luminosity (left panel) and the radio luminosity versus the $H\beta$ broad emission line luminosity (right panel) in our sample. Radio-loud and radio-quiet sources are marked with different symbols to highlight the possible dichotomy of the slope in the figure. It is apparent in Figure 8 that radio-loud and radio-quiet objects almost have the same slope in the left panel (L_X vs. $L_{H\beta}$). The slope is 1.04 ± 0.04 for the radio-quiet subsample and 1.00 ± 0.04 for the radio-loud subsample. The Pearson correlation coefficients are 0.86 and 0.80, respectively. However in the right panel of Figure 8 (L_r vs. $L_{H\beta}$), the slope of radio-loud sources is steeper than that of radio-quiet ones. The slope is 1.68 ± 0.08 for radio-loud sources and 0.99 ± 0.03 for radio-quiet sources, and the Pearson correlation coefficients are 0.64 and 0.88, respectively.

The similar slope of radio-loud and radio-quiet sources in the L_X vs. $L_{H\beta}$ plot seems to indicate that the soft X-ray emission traces well the ionizing luminosity, and is probably isotropic and closely related to the accretion process of the central black hole for both radio-quiet and radio-loud sources. For radio-loud broad-line AGNs, the jet contribution to the soft X-ray emission is probably unimportant. The beaming effect due to the relativistic jet seems also weak in the soft X-ray band. Otherwise, we should expect the large scatter in the L_X vs. $L_{H\beta}$ relation for radio-loud sources in our sample.

The high redshift sources here usually have the measurement of the Mg II $\lambda 2798$ broad emission line instead of the $H\beta$ broad emission line in the observed wavelength. In Figure 9, we show the X-ray luminosity versus the Mg II emission line luminosity in our sample (left panel) and the radio luminosity versus the Mg II emission line luminosity (right panel). The result is consistent with what derived from Figure 8 for low redshift objects.

3.4. Evolution of the black hole fundamental plane

With the currently available large sample, we are able to investigate the evolution of the black hole fundamental plane relation by dividing the sample into different redshift bins. The result is listed in Table 4 and plotted in Figure 10. For radio-quiet objects, the coefficients ξ_{RM} and ξ_{RX} are almost constant in different redshift bins. For radio-loud sources, ξ_{RX} is almost constant with redshift, while ξ_{RM} seems to decrease from positive to negative values as the redshift increases, with the exception of the last redshift bin, where ξ_{RM} is positive again although it shows the largest error bars.

Now we can compare the evolution result obtained by us with some theoretical predic-

tions. From Heinz & Sunyaev (2003), the radio flux produced via the synchrotron radiation from relativistic jet follows the scaling relation

$$F_r \propto M^{\frac{2p+13-(2+p)\alpha_r}{8+2p}} \dot{m}^{\frac{2p+13+(p+6)\alpha_r}{2(p+4)}}. \quad (4)$$

where α_r and p represent the radio spectral index ($f_\nu \propto \nu^{-\alpha_r}$) and the electron energy distribution index ($N \propto E^{-p}$) respectively. M is the central black hole mass. $\dot{m}$ is the dimensionless accretion rate ($\dot{m} = \frac{\dot{M}}{\dot{M}_{Edd}}$). This relation is valid for radio-loud AGNs, whose radio emission is believed to be produced via the synchrotron radiation of the relativistic jet.

The dimensionless accretion rate ($\dot{m}$) is roughly proportional to the ratio between broad emission line and Eddington luminosities ($\frac{L_{\text{broad-line}}}{L_{\text{Edd}}}$) (Wandel, Peterson & Malkan 1999). In the subsection above, we have shown that the soft X-ray luminosity is linearly scaled with the broad emission line luminosity for both radio-loud and radio-quiet subsamples. Therefore, we can use L_X/L_{Edd} to represent the dimensionless accretion rate ($\dot{m}$). After replacing $\dot{m}$ with L_X/L_{Edd} and considering the relation between the Eddington luminosity and the black hole mass ($L_{Edd} \propto M_{BH}$), equation (4) can be turned into

$$L_r \propto M^{-\alpha_r} L_X^{\frac{2p+13+(p+6)\alpha_r}{2(p+4)}}. \quad (5)$$

Therefore, the coefficient ξ_{RM} and ξ_{RX} of the radio-loud subsample can be determined by α_r and p .

In our radio-loud AGN subsample, there are 114 sources that were also detected by the Green Bank 4.85GHz northern sky survey (GB6) (Gregore et al. 1996). Thus we are able to estimate the real radio spectral index α_r ($f_\nu \propto \nu^{-\alpha_r}$) of this small subsample with the observed flux densities at 1.4GHz and 4.8GHz. We plot the radio spectral index versus the redshift for sources in this subsample in Figure 11. There is a weak trend that the spectral index increases with the increasing of the redshift. The mean values of α_r in each redshift bins are -0.28 , -0.12 , -0.31 , -0.27 , -0.06 , -0.11 , and 0.04 (from low- z to high- z). We note that this is probably due to a selection effect in observations, as we may miss high redshift AGNs with the negative spectral index.

Using the average radio spectral indices in different redshift bins, we calculate the theoretical value of the coefficient ξ_{RM} and ξ_{RX} based on equation (5). The general trend of ξ_{RM} is to decrease from 0.28 ($\alpha_r = -0.28$) to -0.04 ($\alpha_r = 0.04$), while ξ_{RX} slightly increases from 1.23 to 1.44. This can be seen clearly from Figure 12. Therefore, the evolution of ξ_{RM} and ξ_{RX} of the radio-loud subsample with the redshift is consistent with the theoretical prediction.

On the other hand, it is also possible that the soft X-ray emission of broad-line AGNs is produced by the synchrotron process in a hot corona around accretion disks. Heinz (2004)

calculated the theoretical correlation between the radio luminosity, X-ray luminosity and black hole mass when the jet radiation dominates. If the physical parameters (i.e., magnetic field strength, electron energy distribution index, etc.) of the hot corona and the relativistic jet are similar, we can use the equations given by Heinz (2004) to roughly estimate the coefficients ξ_{RM} and ξ_{RX} in different redshift bins. We find that with this model, ξ_{RM} decreases from 0.047 ($\alpha_r = -0.28$) to -0.007 ($\alpha_r = 0.04$), and ξ_{RX} increases from 1.23 to 1.44. Therefore, even if the soft X-ray emission mechanism is synchrotron radiation, the observed evolution of the black hole fundamental plane coefficients can still be explained. We can not exclude such a possibility because the origin of the soft X-ray emission is still uncertain.

The radio spectral indices of these 114 radio-loud AGNs obtained from observations allow us to directly calculate their rest-frame 1.4GHz radio fluxes without assuming the canonical value of α_r as 0.5. The black hole fundamental plane relation is fitted based on this small subsample with measured α_r . The derived ξ_{RX} and ξ_{RM} coefficients are 1.34 ± 0.16 and -0.31 ± 0.21 , respectively. These values are consistent within errors with what we obtained from the radio-loud subsample if we set $\alpha_r = 0.5$ ($\xi_{RX} = 1.50 \pm 0.08$ and $\xi_{RM} = -0.20 \pm 0.10$). This indicates that our results for radio-loud AGNs are quite robust.

4. DISCUSSION AND CONCLUSION

4.1. Comparison with previous works

The ξ_{RM} coefficient we derived is rather small, meaning that the dependence of the fundamental plane relation on the black hole mass is weak, which is different from the result obtained by Merloni et al. (2003). However, there are some differences between our sample and that of Merloni et al. (2003). We only include the broad-line AGN, while their sample includes both GBHs and SMBHs. In our sample, the X-ray and radio emission are measured in 0.1-2.4keV and 1.4GHz respectively, while Merloni et al. (2003) used the data of 2-10keV X-ray emission and 5GHz radio core emission. It is still unclear whether the soft (0.1-2.4keV) and hard X-ray (2-10keV) emission have the similar origin for broad-line AGNs. A sample with both available data in the soft and hard X-ray bands will help us address this problem. This is beyond the scope of our current study and will be done in the near future.

Another point worthy to mention is that the black hole mass data in Merloni et al. (2003) were mainly obtained from the literatures. This could introduce scatters due to the different mass measurement techniques. In our current study, the black hole masses of AGNs were estimated from the broad emission line and continuum properties by the same method,

which does not introduce any additional bias. In addition, the black hole masses in our sample span a relatively smaller range than that in Merloni et al. (2003) since we only include broad-line AGNs.

4.2. Relativistic beaming and other uncertainties

Because the radio emission is produced by the relativistic jet, Doppler boosting of the synchrotron radiation (namely the relativistic beaming) would affect the observed radio flux significantly. Here we try to address the question that whether the observed larger fundamental plane coefficient ξ_{RX} (or the larger radio luminosity) of radio-loud sources is due to the Doppler boosting effect.

If the radio emission of both radio-loud and radio-quiet quasars are from jets, the larger radio luminosity observed in radio-loud sources can be considered to have much stronger Doppler boosting effect than radio-quiet AGNs. In other words, we can assume that the radio emission of radio-quiet AGNs is unbeamed. In subsection 3.3, we already show that the soft X-ray emission of broad-line AGNs is probably isotropic and unbeamed for both radio-loud and radio-quiet sources. Therefore, for a radio-loud quasar, its *intrinsic* radio luminosity ($L_{r,jet}$) (unbeamed) may be estimated with its observed X-ray luminosity through the L_r - L_X correlation derived from radio-quiet sources. We will use the ratio between the observed radio luminosity (L_r) and the *intrinsic* radio luminosity ($L_{r,jet}$) to represent the boosting factor of radio-loud sources. The equation of the Doppler boosting effect was given by Heinz & Merloni (2004):

$$L_\nu = \frac{L_{\nu,jet}}{\Gamma^{k+\alpha_r}} \left[\frac{1}{(1 + \beta \cos \theta)^{k+\alpha_r}} + \frac{1}{(1 - \beta \cos \theta)^{k+\alpha_r}} \right]. \quad (6)$$

In Figure 13, we show the distribution of the boosting factor ($L_r/L_{r,jet}$) of the radio-loud subsample (left panel) and the boosting factor as a function of the inclination angle θ (right panel) when different Lorentz factor (Γ) is given. It is apparent in Figure 13 that only with smaller θ ($< 5^\circ$) and larger Γ (> 10), we can produce boosting factors as high as 1000. Such conditions can only be met in BL Lac objects and are not likely to be the case for normal broad-line AGNs studied in this work. With the typical Lorentz factor of $\Gamma \sim 5$ for broad-line AGNs (Orr & Browne 1982) and a non-negligible inclination angle ($\theta \gtrsim 10^\circ$) (Maraschi et al. 1994), the boosting value is estimated to be less than 30. As can be seen in Figure 13, there are about half of radio-loud sources whose boosting factors are larger than the predicted maximum boosting value. Therefore, although the Doppler boosting effect

indeed has significant influence on the radio emission of radio-loud AGNs, this effect alone is not enough to explain the observed larger radio luminosity of radio-loud broad-line AGNs. This is consistent with the result given by Heinz & Merloni (2004) for the case of unbeamed X-ray emission.

The non-simultaneous observations in the radio, X-ray and optical bands for sources in our sample may lead to other uncertainties. Both the ROSAT and FIRST surveys were conducted in the 1990s (Becker et al. 1995; Voges et al. 1999; Britzen et al. 2007). SDSS-I observations started from 1998, and ended in 2005 (York et al. 2000). So the data in different bands were obtained within ten years or so. Unless the luminosities of most objects in our sample varied significantly in these years, our result may not be affected too much due to this effect. In addition, several (2~3) factors change of the luminosities of some individual objects have little influence on the statistically significant results derived here for a large sample. However, as broad-line AGNs often show X-ray variabilities in timescales from hours to decades, the non-simultaneous observations could be an issue if the X-ray fluxes of AGNs vary significantly. Therefore, simultaneous observations in different bands for a sample of broad-line AGNs are still needed to confirm our results, although they are difficult to conduct for a large sample.

4.3. Emission mechanisms

The existence of the AGN radio-loud and radio-quiet dichotomy is still an unsolved issue. There are evidences that the radio emission of radio-quiet AGNs is likely produced by a weak (sub-relativistic) jet near the black hole (Blundell & Beasley 1998; Leipski et al. 2006), while radio-loud quasars are usually associated with large scale jets of higher radio power (Rawlings & Saunders 1991; Miller, Rawlings & Saunders 1993). Here we fit the black hole fundamental plane relation for a broad-line AGN sample and find that the coefficients are quite different between radio-loud and radio-quiet quasars, especially the coefficient ξ_{RX} , which is steeper for radio-loud objects. We also find that the larger radio luminosities of radio-loud AGNs can not be produced by the Doppler boosting effect alone. Therefore, the radio emission mechanism may be quite different between radio-loud and radio-quiet AGNs. Such kind of difference can be caused by many uncertain physical parameters, such as the detailed disk/jet magnetic field strength and configuration, the electron energy distribution, and the black hole spin, etc.(Heinz & Sunyaev 2003; Sikora et al. 2007). More detailed observational and theoretical studies are still required.

For the soft X-ray emission, we have shown that the correlation slopes between the soft X-ray luminosity and the broad emission line luminosity are all around 1 for both radio-

loud and radio-quiet AGNs. Therefore, the soft X-ray emission is probably produced via the accretion process near the central black hole, and it is unbeamed and isotropic for both radio-loud and radio-quiet sources. Otherwise larger scatters should exist in the L_X vs. $L_{H\beta}$ (L_{MgII}) relation. However, there are many uncertain factors related to the origin of the soft X-ray emissions of AGNs (i.e., the accretion flow, hot corona, and warm absorber, etc.). A detailed study of the soft X-ray spectra for a larger sample of broad-line AGNs may give us clues to understand the origin of the soft X-ray emissions.

4.4. Conclusion

We revisited the fundamental plane relation of the black hole activity based on a large broad-line AGN sample selected on the basis of the cross-identification of the RASS, SDSS, and FIRST catalogs. The results of our work confirm the main result of Wang et al. (2006), namely, the black hole fundamental plane relation of the radio-quiet subsample is different from that of the radio-loud subsample; the coefficient ξ_{RX} becomes larger as the radio-loudness increases; the black hole mass seems unimportant in the black hole fundamental plane relation. We also found that the soft X-ray emission is most likely produced via the accretion process of the central black hole for both radio-quiet and radio-loud sources. In particular, for radio-loud sources, the jet contribution to the soft X-ray emission seems unimportant and the Doppler boosting of the relativistic jet is also weak in the soft X-ray band. Moreover, by dividing the radio-loud and radio-quiet samples into different redshift bins, we studied the evolution of the fundamental plane relation. For radio-quiet sources, there seems to be no clear evolution, while for radio-loud objects, the correlation coefficient ξ_{RM} tends to decrease as the redshift increases. We found that the evolution of the radio spectral index can help us at least partly understand such an evolution. Finally, we briefly discussed the beaming effect and some other uncertainties associated with the fundamental plane relation derived here. We found that Doppler boosting effect indeed has significant influence on the radio emission of radio-loud AGNs, but this effect alone is not enough to explain the observed radio luminosity of the radio-loud sources.

We thank Fukun Liu, Lei Qian, Xian Chen, Shuo Li, Da-Wei Xu and Weimin Yuan for helpful discussions, and Eric Peng for checking the English. We are also grateful to the anonymous referee for his/her helpful comments. This work is supported by the NSFC grant No.10525113, the RFDP grant 20050001026, the NCET grant (NCET-04-0022) and the 973 Program No.2007CB815405. Funding for the SDSS has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese

Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS Web Site is <http://www.sdss.org/>. The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.

REFERENCES

- Akritas, M. G., & Siebert, J. 1996, MNRAS, 278, 919
- Anderson, S. F. et al. 2003, AJ, 126, 2009
- Anderson, S. F. et al. 2007, AJ, 133, 313
- Avni, Y., & Tananbaum, H. 1982, ApJ, 262, L17
- Becker, R. H., Helfand, D. J., White, R. L., Gregg, M. D., & Laurent-Muehleisen, S. A. 2003, VizieR Online Data Catalog, 8071
- Becker, R. H., White, R. L., & Helfand, D. J. 1995, ApJ, 450, 559
- Begelman, M. C., Blandford, R. D., & Rees, M. J., 1984, Rev. Mod. Phys. 56, 255
- Blundell, K. M., & Beasley, A. J. 1998, MNRAS, 299, 165
- Boroson, T. A., & Green, R. F. 1992, ApJS, 80, 109
- Bregman, J. N. 2005, ApJ, astro-ph/0511368
- Brinkmann W., Laurent-Muehleisen S. A., Voges W., Siebert J., Becker R. H., Brotherton M. S., White R. L., Gregg M. D. 2000, A&A, 356, 445
- Brinkmann W., Yuan W., & Siebert J., 1997, A&A 319, 413
- Britzen, S. et al. 2007, A&A, 476, 759
- Canosa, C.M., Worrall, D.M., Hardcastle, M.J., & Birkinshaw, M. 1999, MNRAS, 310, 30
- Colin, S., & Kawaguchi, T. 2004, A&A, 426, 797
- Falcke, H., Körding, E., & Markoff, S. 2004, A&A, 414, 895
- Fender, R. P., Gallo, E., & Jonker, P. G. 2003, MNRAS, 343, L99
- Frank, J., King, A., & Raine, D., 2002, Accretion Power in Astrophysics, 3rd Edition, Cambridge University Press
- Gallo, E., Fender, R. P., & Pooley, G. G. 2003, MNRAS, 344, 60
- Gallo, E., 2006, AIP Conference Proceedings, Vol. 924, 715
- Gregory, P. C., Scott, W. K., Douglas, K., & Condon, J. J. 1996, ApJS, 103, 427

- Haardt, F. & Maraschi, L. 1993, *ApJ*, 413, 507
- Heinz, S., & Merloni, A. 2004, *MNRAS*, 355, L1
- Heinz, S., & Sunyaev, R. A. 2003, *MNRAS*, 343, L59
- Heinz, S. 2004, *MNRAS*, 355, 835
- Isobe, T., Feigelson, E.D., Akritas, M.G. & Babu, G.J. 1990, *ApJ*, 364, 104
- Kaspi, S., Smith, P. S., Netzer, H., Peterson, B. M., Vestergaard, M., & Giveon, U. 2000, *ApJ*, 533, 631
- Kaspi, S., Maoz, D., Netzer, H., Peterson, B. M., Vestergaard, M., & Jannuzi, B. T. 2005 *ApJ*, 629, 61
- Kellermann, K. I., Sramek, R., Schmidt, M., Shaffer, D. B., & Green, R. 1989, *AJ*, 98, 1195
- Komossa, S., Voges, W., Xu, D., Mathur, S., Adorf, H. M., Lemson, G., Duschl, W. J., & Grupe, D. 2006, *AJ*, 132, 531
- Körding, E., Falcke H. & Corbel S. 2006, *A&A*, 456, 439
- Lacy, M., Laurent-Muehleisen, S. A., Ridgway, S. E., Becker, R. H., & White, R. L. 2001, *ApJ*, 551L, 17L
- Leipski, C., Falcke, H., Bennert, N., & Huttemeister, S. 2006, *A&A*, 455, 161
- Maraschi, L., & Rovetti, F. 1994, *ApJ*, 436, 79
- McLure, R. J., & Jarvis, M. J. 2002, *MNRAS*, 337, 109
- Merloni, A., Heinz, S., & Di Matteo, T. 2003 *MNRAS*, 345, 165
- Merloni, A., Kording, E., Heinz, S., Markoff, S., DiMatteo, T., & Falcke, H. 2006, *New Astronomy* 11, 567
- Miller, P., Rawlings, S., & Saunders, R. 1993, *MNRAS*, 263, 425
- Orr, M. J. L., & Browne, I. W. A. 1982, *MNRAS*, 200, 1067
- Osterbrock, D. E., & Pogge, R.W. 1985, *ApJ*, 297, 166
- Panessa, F., Barcons, X., Bassani, L., Cappi, M., Carrera, F. J., Ho, L. C., & Pellegrini, S. 2007, *A&A*, 467, 519

- Rawlings, S., & Saunders, R. 1991, *Nature*, 349, 138
- Shakura, N. I. & Sunyaev, R.A. 1973, *A&A*, 24, 337
- Sikora, M., Stawarz, L., & Lasota, J. P. 2007, *ApJ*, 658, 815
- Voges, W. et al. 1999, *A&A*, 349, 389
- Wandel, A., Peterson, B. M., & Malkan, M. A. 1999, *ApJ*, 526, 579
- Wang, R., Wu, X-B., Kong, M. Z. 2006, *ApJ*, 645, 890
- White, R. L., Becker, R. H., Helfand, D. J., & Gregg, M. D. 1997, *ApJ*, 475, 479
- White, R. L. et al. 2000, *ApJS*, 126, 133
- Wu, X-B., Wang, R., Kong, M. Z., Liu, F. K., & Han, J. L. 2004, *A&A*, 424, 793
- Xue, Y. Q., & Cui, W. 2007, *A&A*, 466, 1053
- Xue, Y. Q., Wu, X.-B., & Cui, W. 2008, *MNRAS*, 384, 440
- York, D. G. et al. 2000, *AJ*, 120, 1579
- Yuan, F., & Cui, W. 2005, *ApJ*, 629, 408
- Yuan, W., Zhou, H. Y., Komossa, S., Dong, X. B., Wang, T. G., Lu, H. L., & Bai, J. M. 2008, *ApJ*, in press (astro-ph/0806.3755)

A. RADIO-LOUD NARROW-LINE SEYFERT 1 GALAXIES IN OUR SAMPLE

Narrow-Line Seyfert 1 (NLS1) galaxies are a sub-class of AGN population. Their optical broad permitted emission lines are usually narrower ($\text{FWHM}_{\text{H}\beta} < 2000 \text{ km/s}$) than that in normal Broad-Line Seyfert 1 (BLS1) galaxies (Osterbrock & Pogge 1985). The NLS1s also show weak $[\text{OIII}]\lambda 5007/\text{H}\beta_{\text{total}}$ emission and strong FeII emissions (Boroson & Green 1992). Recently, Komossa et al. (2006) argued that the classical criteria ($\text{FWHM}_{\text{H}\beta} < 2000 \text{ km/s}$) is not well defined or even completely arbitrary. They suggested that $R_{4570} > 0.5$ may be a physically more meaningful criteria to distinguish the NLS1 from the ordinary BLS1.⁶

The study of the physical mechanism of NLS1s is still ongoing. There is growing evidence that most NLS1s are objects with low black hole masses and high accretion rates, close to or even above the Eddington accretion rate (Collin & Kawaguchi 2004).

Radio-loud NLS1s are rare objects in NLS1 population (Komossa et al. 2006). In order to understand their radio properties, it is important to expand the number of radio-loud NLS1s. Anderson et al. (2007) roughly examined the optical broad permitted emission line of AGNs in their catalog. They marked those objects that have $\text{FWHM}_{\text{H}\beta} < 2000 \text{ km/s}$ as “NLS1?” in the comment columns of their tables. Seventy-four of these objects have optical spectra from SDSS and are detected in the FIRST radio survey, and thus are included in our analysis in this paper. We calculated the radio-loudness values for these objects, and found that five of them are radio-loud. We list the properties of the five sources in Table A1.

In a newly published work, Yuan et al. (2008) present a comprehensive study of a sample of 23 radio-loud NLS1 galaxies. Among those 23 sources, two radio-loud NLS1s are also discovered in our present work, which are SDSS J144318.56+472556.7 and SDSS J114654.28+323652.3. These two independent studies confirm the nature of radio-loud NLS1s of these two objects.

⁶The optical Fe II strength, R_{4570} , is the ratio of the Fe II complex between the rest wavelength $\lambda 4434\text{\AA}$ and $\lambda 4684\text{\AA}$ to the total $\text{H}\beta$ flux, including the narrow component (Boroson & Green 1992)

Table 1. The AGN Samples^a

Name	z	$\log(L_X/1 \text{ erg s}^{-1})$	$\log(L_r/1 \text{ erg s}^{-1})$	log R	$\log(M/M_\odot)$	$L_{\text{broad-line}}$ ^b	flag ^c
SDSS J000608.04 – 010700.7	0.949	45.60	41.13	1.345	8.71	43.47	0
SDSS J000710.01 + 005329.1	0.316	44.95	39.79	0.549	9.07	42.90	1
SDSS J004319.73 + 005115.4	0.308	44.64	39.61	0.506	9.42	42.90	1

^aOnly 3 rows of the catalog are shown here. A complete catalog will be provided in the electronic version

^bThe broad emission line luminosity. It is represented by the H_β broad component luminosity or the Mg II emission line luminosity (when H_β is unavailable).

^c1 means that L_{H_β} is used for $L_{\text{broad-line}}$, 0 means that L_{MgII} is used for $L_{\text{broad-line}}$.

Table 2. Derived Fundamental Plane Relation

Subsample	Number	ξ_{RX}	ξ_{RM}	Constant	σ_r
Total	725	1.47 ± 0.06	0.04 ± 0.07	-0.33 ± 0.06	0.83
Radio-quiet	227	0.73 ± 0.10	0.31 ± 0.12	-0.68 ± 0.07	0.42
Radio-loud	498	1.50 ± 0.08	-0.20 ± 0.10	0.05 ± 0.10	0.75
Merloni et al. (2003)	...	0.60 ± 0.11	$0.78^{+0.11}_{-0.09}$	$7.33^{+4.05}_{-4.07}$	0.88
Total ^a	115	1.33 ± 0.15	0.30 ± 0.18	-0.40 ± 0.14	0.89
Radio-quiet ^a	39	0.85 ± 0.10	0.12 ± 0.13	-0.77 ± 0.07	0.38
Radio-loud ^a	76	1.39 ± 0.17	0.17 ± 0.21	-0.17 ± 0.21	0.77

^aResults derived by Wang et al. (2006)

Table 3. Partial Correlation Test For The Fundamental Plane Correlation

X	Y	Z	Type	Number	τ	σ	P_{null}
log Lx	log Lr	log D	Radio-loud	498	0.321	0.0264	<1.000E-10
log Lx	log Lr	log D	Radio-quiet	227	0.270	0.0424	1.916E-10
log Lx	log Lr	log D	Total	725	0.308	0.0220	<1.000E-10
log Lx	log Lr	log M	Radio-loud	498	0.567	0.0201	<1.000E-10
log Lx	log Lr	log M	Radio-quiet	227	0.578	0.0366	<1.000E-10
log Lx	log Lr	log M	Total	725	0.598	0.0188	<1.000E-10
log(Lx/Ledd)	log(Lr/Ledd)	log D	Radio-loud	498	0.472	0.0249	<1.000E-10
log(Lx/Ledd)	log(Lr/Ledd)	log D	Radio-quiet	227	0.517	0.0340	<1.000E-10
log(Lx/Ledd)	log(Lr/Ledd)	log D	Total	725	0.412	0.0197	<1.000E-10
log(Lx/Ledd)	log(Lr/Ledd)	log M	Radio-loud	498	0.572	0.0204	<1.000E-10
log(Lx/Ledd)	log(Lr/Ledd)	log M	Radio-quiet	227	0.553	0.0325	<1.000E-10
log(Lx/Ledd)	log(Lr/Ledd)	log M	Total	725	0.550	0.0160	<1.000E-10

Table 4. Derived Fundamental Plane Relation In Different Redshift Bins

Redshift Range	ξ_{RX}	ξ_{RM}	N
Radio-Quiet subsample			
$z < 0.13$	0.33 ± 0.20	0.23 ± 0.19	77
$0.13 < z < 0.4$	0.35 ± 0.23	0.16 ± 0.20	81
$0.4 < z < 1.0$	0.49 ± 0.34	0.12 ± 0.34	48
$z > 1.0$	0.55 ± 0.69	0.45 ± 0.77	21
Radio-Loud subsample			
$z < 0.3$	0.63 ± 0.24	0.15 ± 0.23	58
$0.3 < z < 0.45$	1.59 ± 0.38	-0.26 ± 0.24	67
$0.45 < z < 0.65$	0.81 ± 0.35	-0.03 ± 0.21	86
$0.65 < z < 0.80$	1.73 ± 0.53	-0.21 ± 0.24	71
$0.80 < z < 1.0$	0.70 ± 0.40	-0.91 ± 0.41	64
$1.0 < z < 1.3$	0.95 ± 0.44	-0.66 ± 0.34	77
$z > 1.3$	1.27 ± 0.38	0.11 ± 0.37	75

Table A1. Radio-Loud Narrow Line Seyfert 1 Objects

Name	z	$\text{FWHM}_{H\beta}/1 \text{ km s}^{-1}$	$\log(M_{\text{BH}}/M_{\odot})$	$\log R$	R_{4570}
SDSS J144318.56 + 472556.7 ^a	0.703	1810.1	7.14	2.91	5.50
SDSS J114654.28 + 323652.3	0.465	2374.3	7.43	1.98	1.59
SDSS J073320.84 + 390505.2	0.664	2867.8	7.86	2.80	3.27
SDSS J154510.96 + 345246.9	0.516	3269.7	8.13	1.34	0.74
SDSS J123304.05 – 003134.1 ^a	0.471	3297.7	7.81	2.32	2.18

^a H_{β} emission line is fitted with just one Gaussian




Fig. 1.— Global properties of our AGN sample, with the top panels showing the histograms of the redshift and logarithm of radio-loudness, the middle panels showing the histograms of the X-ray luminosity and black hole mass (in units of $M_{\odot}$), and the bottom panels showing the radio luminosity in logarithm units and ratios of X-ray to Eddington luminosity.




Fig. 2.— Rest-frame 1.4GHz radio luminosity (L_r) vs. the 0.1-2.4 keV X-ray luminosity, with different symbols representing different logarithmic bins of the black hole mass (in units of $M_\odot$). In the left panel we plot the logarithm of the luminosity directly, while in the right panel we scale the radio and X-ray luminosity with L_{Edd} .




Fig. 3.— Rest-frame 1.4GHz radio luminosity (L_r) vs. the 0.1-2.4 keV X-ray luminosity, with different symbols representing different radio-loudness bins. We plot the radio vs. X-ray luminosity in the left panel, and the Eddington luminosity scaled radio vs. X-ray luminosity in the right panel.




Fig. 4.— Ratio of the radio to X-ray luminosity vs. the black hole mass. Different symbols represent objects in different radio-loudness bins.




Fig. 5.— Edge-on view of the black hole fundamental plane for all AGNs in our sample. Different symbols represent sources in different radio-loudness bins. The dashed line is the best-fitting result for radio-quiet AGNs.




Fig. 6.— Correlation between $\log(L_r/L_{Edd})$ and $0.96 \times \log(L_X/L_{Edd})$. The symbols and lines have the same meanings as in Figure 5.




Fig. 7.— Results of the Monte Carlo test using scrambled radio fluxes. The position of the vertical line represents Pearson’s correlation coefficient of the real dataset.




Fig. 8.— Plot of the 0.1-2.4keV X-ray luminosity versus the $H\beta$ broad emission line luminosity (left panel) and the rest-frame 1.4GHz radio luminosity versus the $H\beta$ broad emission line luminosity (right panel). The meanings of symbols and lines are: green diamond: radio-loud AGNs; black cross: radio-quiet AGNs; black line: best-fitting result for the radio-quiet subsample; dashed black line: best-fitting result for the radio-loud subsample.




Fig. 9.— Plot of the 0.1-2.4keV X-ray luminosity versus the Mg II emission line luminosity (left panel) and the rest-frame 1.4GHz radio luminosity versus the Mg II emission line luminosity (right panel). The symbols and lines have the same meanings as in Figure 8.




Fig. 10.— Dependence of the black hole fundamental plane coefficients ξ_{RM} and ξ_{RX} on redshift for radio-quiet and radio-loud AGNs.




Fig. 11.— Redshift dependence of the radio spectral index α_r of the 114 radio-loud AGNs detected both at 1.4GHz and 5GHz frequencies.




Fig. 12.— Predicted black hole fundamental plane coefficients based on equation (5). Here, α_r is the radio spectral index and p equals to 2.




Fig. 13.— The histogram of the boosting factor ($L_r/L_{r,jet}$) of the radio-loud subsample (left panel) and the boosting factor as a function of the inclination angle θ (right panel), with different line representing different Lorentz factor (Γ).